

MATH 401: Lecture 2 (08/21/2025)

Today: *sets and operations

Sets and Operations (LS IRA 1.2)

Set: Collection of mathematical objects.

They can be finite, e.g., $\{2, 5, 9, 1, 6\}$, or infinite, e.g., $[0, 1]$, the collection of all $x \in \mathbb{R}$ with $0 \leq x \leq 1$.

→ "element of"

→ set of all real numbers

Given sets A, B we have

$A \subseteq B$: A is a subset of, or equal to, B .

$A \subset B$: A is a strict subset of B , i.e., there is at least one $x \in B$ such that $x \notin A$.

But $\nexists x \in A, x \in B$ holds.

To prove $A = B$, we often prove $A \subseteq B$ and $A \supseteq B$ ($\text{or } B \subseteq A$).

Here are some standard sets we will use regularly.

\emptyset : empty set.

$\mathbb{N} = \{1, 2, 3, \dots\}$, set of all natural numbers

\mathbb{R} = set of all real numbers

$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$, set of all integers

\mathbb{Q} = set of rational numbers, \mathbb{C} = set of complex numbers.

\mathbb{R}^n : set of all real n -tuples, or n -vectors

Notation for sets: $[-2, 1] = \{x \in \mathbb{R} \mid -2 \leq x \leq 1\}$.

closed interval from -2 to 1

→ "such that"

could also use ":" instead of "1".

More generally, $A = \{a \in B \mid P(a)\}$.

↓
bigger set
than A

property

Union and Intersection

If A_i are sets for $i=1, \dots, n$, i.e., A_1, A_2, \dots, A_n are sets, then



$\bigcup_{i=1}^n A_i = A_1 \cup A_2 \cup \dots \cup A_n = \{a \mid a \in A_i \text{ for at least one } i\}$ is their union,
 ↗ "for all"

$\bigcap_{i=1}^n A_i = A_1 \cap A_2 \cap \dots \cap A_n = \{a \mid a \in A_i \forall i\}$ is their intersection.

universe $\rightarrow \bigcup$

$A \cup B$


$A \cap B$



LSIRA 1.2 Prob 1 Show $[0, 2] \cup [1, 3] = [0, 3]$.

We show $[0, 2] \cup [1, 3] \subseteq [0, 3]$ and

$[0, 2] \cup [1, 3] \supseteq [0, 3]$.



(\subseteq) Let $x \in [0, 2] \cup [1, 3]$

$\Rightarrow x \in [0, 2]$ or $x \in [1, 3]$ (definition of \cup).

$x \in [0, 2] \Rightarrow x \in [0, 3]$ (as $[0, 3]$ contains $[0, 2]$)

$x \in [1, 3] \Rightarrow x \in [0, 3]$. In either case, $x \in [0, 3]$.

Hence $[0, 2] \cup [1, 3] \subseteq [0, 3]$.

(\supseteq) Let $x \in [0, 3]$. Hence $0 \leq x \leq 3$. Then we get that

either $x \leq 2$, and hence $x \in [0, 2]$, or $x \in (2, 3]$.

But if $x \in (2, 3]$ then $x \in [1, 3]$ (as $[1, 3]$ includes $(2, 3]$).

$\Rightarrow x \in [0, 2] \cup [1, 3]$.

Hence $[0, 3] \subseteq [0, 2] \cup [1, 3]$.

The result is an obvious one. But we go through the steps of a formal proof more for practice!

□

Distributive Laws of Union and Intersection

For all sets B, A_1, \dots, A_n , we have

$$\text{LSIRA} \quad (1.2.1) \quad B \cap (A_1 \cup A_2 \cup \dots \cup A_n) = (B \cap A_1) \cup (B \cap A_2) \cup \dots \cup (B \cap A_n).$$

Using more compact notation, we can write

$$B \cap \left(\bigcup_{i=1}^n A_i \right) = \bigcup_{i=1}^n (B \cap A_i)$$

Proof

We will prove

$$B \cap (A_1 \cup \dots \cup A_n) \subseteq (B \cap A_1) \cup \dots \cup (B \cap A_n), \text{ and}$$

$$B \cap (A_1 \cup \dots \cup A_n) \supseteq (B \cap A_1) \cup \dots \cup (B \cap A_n).$$

(\subseteq) Let $x \in B \cap (A_1 \cup \dots \cup A_n)$.

$$\Rightarrow x \in B \text{ and } x \in (A_1 \cup \dots \cup A_n) \quad (\text{definition of } \cap)$$

$$\Rightarrow x \in B \text{ and } x \in A_i \text{ for at least one } A_i. \quad (\text{defn. of } \cup)$$

$$\Rightarrow x \in B \cap A_i \text{ for at least one } A_i.$$

$$\Rightarrow x \in (B \cap A_1) \cup \dots \cup (B \cap A_n).$$

(\supseteq) Let $x \in (B \cap A_1) \cup \dots \cup (B \cap A_n)$.

$$\Rightarrow x \in (B \cap A_i) \text{ for at least one } A_i.$$

$$\Rightarrow x \in B \text{ and } \underbrace{x \in A_i \text{ for at least one } A_i}_{\text{for at least one } A_i}$$

$$\Rightarrow x \in B \text{ and } x \in (A_1 \cup \dots \cup A_n)$$

$$\Rightarrow x \in B \cap (A_1 \cup \dots \cup A_n).$$

LSIRA (1.2.2) is assigned in Homework 1. □

Set Difference and Complement

We write $A \setminus B$ or $A - B$ "setminus"


Caution!

* $A \setminus B \neq B \setminus A$!

" A setminus B " is $A \setminus B = \{a \mid a \in A, a \notin B\}$.

If U is the universe, i.e., $A \subseteq U$ for all sets A , then

$A^c = U \setminus A = \{a \in U \mid a \notin A\}$ is the complement of A (or A -complement).



De Morgan's Laws

LSIRA
(1.2.3)

$$(A_1 \cup \dots \cup A_n)^c = A_1^c \cap \dots \cap A_n^c.$$

"complement of union = intersection of complements"


LSIRA
(1.2.4)

$$(A_1 \cap \dots \cap A_n)^c = A_1^c \cup A_2^c \cup \dots \cup A_n^c.$$

complement of intersection = union of complements.

See LSIRA for the proof.

Let's illustrate (1.2.4) for $n=2$, i.e., with A_1 and A_2 first.



We will prove subset inclusion in both directions.

(\subseteq) Let $x \in (A_1 \cap \dots \cap A_n)^c$

$$\Rightarrow x \notin A_1 \cap \dots \cap A_n \quad (\text{definition of complement})$$

$$\Rightarrow x \notin A_j \text{ for some } j. \quad (\text{definition of } \cap)$$

$$\Rightarrow x \in A_j^c \text{ for some } j$$

$$\Rightarrow x \in A_1^c \cup \dots \cup A_n^c.$$

$$\text{Hence } (A_1 \cap \dots \cap A_n)^c \subseteq A_1^c \cup \dots \cup A_n^c.$$

(\supseteq) Let $x \in A_1^c \cup \dots \cup A_n^c$.

$$\Rightarrow x \in A_j^c \text{ for some } j$$

$$\Rightarrow x \notin A_j \text{ for some } j$$

$$\Rightarrow x \notin A_1 \cap \dots \cap A_n.$$

$$\Rightarrow x \in (A_1 \cap \dots \cap A_n)^c.$$

since $x \notin A_j$ for some j , it cannot be in the intersection of all A_i 's.

$$\text{Hence } A_1^c \cup \dots \cup A_n^c \subseteq (A_1 \cap \dots \cap A_n)^c.$$

□

Cartesian Products

For A, B : sets, we define

$$A \times B = \{(a, b) \mid a \in A, b \in B\}$$

→ cartesian product of A and B

Given $A_i, i=1, \dots, n$ (A_1, \dots, A_n), we define

$$A_1 \times A_2 \times \dots \times A_n = \prod_{i=1}^n A_i = \{(a_1, \dots, a_n) \mid \underbrace{a_i \in A_i}_{a_1 \in A_1, a_2 \in A_2, \dots, a_n \in A_n} \forall i\}$$

→ compact notation
 \prod : product

$$A_1 \times A_2 \times \dots \times A_n = \prod_{i=1}^n A_i = \{(a_1, \dots, a_n) \mid \underbrace{a_i \in A_i}_{a_1 \in A_1, a_2 \in A_2, \dots, a_n \in A_n} \forall i\}$$

e.g., \mathbb{R}^n : set of n -tuples of real numbers
 (or set of real n -vectors)

LSIRAI.2 Prob 9 (Pg 11) Prove that $(A \cup B) \times C = (A \times C) \cup (B \times C)$.

We'll finish the proof in the next lecture..